



# Pharmacy Friday

Brief pearls related to acute care pharmacology and evidence-based medicine

Other pearls found at:

- <https://sites.google.com/presby.edu/pharmacy-friday>



## Airway Series: Induction Medications

### Introduction

1. Rapid sequence intubation (RSI) is a process whereby an induction agent and a neuromuscular blocking agent are given in rapid succession to facilitate endotracheal intubation
2. The selection of a specific sedative depends on multiple factors: the clinical scenario, which includes patient factors (includes cardiorespiratory and neurologic status, allergies, comorbidity) and the clinician's experience/training and institutional factors, as well as the characteristics of the sedative
3. Etomidate remains the most commonly used induction agent, however, it is not without its own pharmacologic considerations
4. The use of ketamine is continuing to rise especially due to its unique pharmacologic profile and its niche is becoming prevalent in situations where the risk of hypotension is significant

Pharmacology			
	Etomidate	Ketamine	Propofol
<b>Dose</b>	0.3 mg/kg IV	1-2 mg/kg	1.5-2 mg/kg
<b>Administration</b>	IV push	IV Push	IV push
<b>Formulation</b>	20 mg/ 10 ml vial	Prefilled 50 mg/5 ml Syringe	1000 mg/100 ml vial
<b>PK/PD</b>	Onset: ~20 seconds Duration: 4-10 minutes Metabolism: Hydrolysis of the ethylester side Renal Excretion: 75%	Onset: ~ IV 30 seconds IM 3-4 minutes Duration: 5-10 minutes Metabolism: N- demethylation Renal Excretion: 91%	Onset: ~10-50 seconds Duration: 3-10 minutes Metabolism: CYP2B6 Renal Excretion: 88%
<b>Adverse Effects</b>	Injection site pain, nausea, vomiting, myoclonus	Hypertension, tachycardia, emergency phenomenon	Hypotension, bradycardia
<b>Drug Interactions</b>	No major reactions	No major reactions	No major reactions
<b>Compatibility</b>	Incompatible with vitamin c and vecuronium	Incompatible with furosemide, insulin, phenytoin, and sodium bicarbonate	Incompatible with methylprednisolone, phenytoin, and metoclopramide
<b>Location in GHS</b>	CPR, Trauma room, Zone 2+3	CPR, Trauma room, Zone 2+3	CPR, Trauma room, Zone 2+3
<b>Comments</b>	There is hypothetical concerns about adrenal insufficiency with a single dose. <b>Hemodynamically neutral</b>	Rapid IV push may cause apnea, Option for delayed sequence intubation. <b>Increase BP and HR</b>	Large dose rapid doses can cause large <b>drops in HR and BP. Option for increase ICP</b>

Drug	Hemodynamic Effect	Comments
<b>Etomidate</b>	↔ BP, ↔ CO, ↔ HR, ↓ cortisol, ↔ ICP	Prolonged inhibition of steroid synthesis in the critically ill; withdrawn from number of countries
<b>Ketamine</b>	↑BP, ↑ HR, ↑ CO, ↔ cortisol, ↑↓ ICP	↔ or ↑ CPP and ↔ ICP with standard anesthetic management
<b>Propofol</b>	↓ BP, ↔ HR, ↓ CO, ↔ cortisol, ↓ ICP	Hemodynamic compromise marked in elderly, ASA 3 or more or hypovolemic patients with 'standard' induction dose

## Comments by ED Physician Attendings

	Pros	Cons
<b>Ketamine</b>	Ketamine has some bronchodilatory properties and can be useful if intubating for asthma angioedema, airway narrowing from anaphylaxis, infection or malignant processes are the typical examples	The dose should be greatly reduced in shock states- most notably hypovolemic shock as it is a direct myocardial depressant  There are some case reports of cardiac arrest when full induction doses of ketamine are pushed in these patients. In those patients I will push 10mg at a time until dissociation occurs (usually around 0.2-0.3 mg/kg in my experience).
<b>Etomidate</b>	Most commonly used unless there circumstance where the patient will not be paralyzed such as difficult airways such as angioedema, airway narrowing from anaphylaxis, infection or malignant processes	Very short duration of action is important- 3 to 5 minutes  Etomidate with rocuronium can be a recipe for paralysis without sedation unless you are right on top of providing post intubation sedation
<b>Propofol</b>	Due to vasodilatory and anti-epileptic properties, propofol is most useful in hypertensive head bleeds and patients with status epilepticus those with enough BP to work with but titrate 10 mg at a time	Hypotension and bradycardia should be noted, especially in trauma patients

### Overview of Evidence

Author, year	Design/ sample size	Intervention & Comparison	Outcome
Dietrich, 2018	Retrospective review/ n=83	Propofol vs Non-propofol (etomidate or midazolam)	↑ <b>post-intubation hypotension</b> with propofol OR 3.64 (95% CI 1.16-13.24)  <b>Similar rates of hypotension</b> were seen among patients who received ≤2 mg/kg and those receiving >2 mg/kg  No significant differences between groups in <b>hospital length of stay or mortality</b>
Lyons, 2015	Cohort study/ n=261	Etomidate+ Succinylcholine (Group 1) vs Fentanyl+ ketamine+ rocuronium (Group 2)	Significantly better laryngeal views with fentanyl/ketamine/rocuronium group  <b>100% first attempt intubation</b> with fentanyl/ketamine/rocuronium group  ↑ <b>post-intubation MAP+ HR</b> with etomidate + succinylcholine
Bruder, 2015	Cochrane Review	Etomidate Midazolam Propofol Ketamine	There was <b>no difference</b> in mortality, hospital LOS, duration of ventilation, and duration of vasopressors  Etomidate associated with ↑ <b>ACTH</b> and ↓ <b>in cortisol level</b>
Tekwani K, 2010	RCT/ n=122	Etomidate 0.3 mg/kg vs midazolam 0.1 mg/kg	<b>No significant differences</b> in median hospital LOS (9.5 vs 7.3 days), ICU LOS (4.2 vs 3.1 days), In-hospital mortality ( 26% vs 43%) or ventilator days
Jabre P, 2009	RCT/ n=469	Etomidate 0.3 mg/kg vs Ketamine 2 mg/kg	<b>No difference in intubating condition</b> , SOFA score, 28 day mortality, Vent free days, vasopressor support, or GCS
White, 1982	RCT/ n= 80	Ketamine 1.5 mg/kg  Thiopental 4 mg/kg  Midazolam 0.3 mg/kg  Midazolam 0.15 mg/kg + ketamine 0.75 mg/kg	Thiopental ↓ MAP by 11%, ketamine increased MAP by 10%, while neither midazolam nor the midazolam-ketamine combination significantly changed MAP  Midazolam effectively attenuated both the cardiostimulatory responses and unpleasant emergence reactions associated with ketamine

### References

1. Micromedex [Electronic version]. Greenwood Village, CO: Truven Health Analytics. Retrieved September 6, 2018, from <http://www.micromedexsolutions.com/>
2. Dietrich SK. Hemodynamic Effects of Propofol for Induction of Rapid Sequence Intubation in Traumatically Injured Patients. Am Surg. 2018 Sep 1;84(9):1504-1508.
3. White PF. Comparative evaluation of intravenous agents for rapid sequence induction—thiopental, ketamine, and midazolam. Anesthesiology. 1982 Oct;57(4):279-84.
4. Jabre P. Etomidate versus ketamine for rapid sequence intubation in acutely ill patients: a multicentre randomised controlled trial. Lancet. 2009 Jul 25;374(9686):293-300.
5. Tekwani KL. A comparison of the effects of etomidate and midazolam on hospital length of stay in patients with suspected sepsis: a prospective, randomized study. Ann Emerg Med. 2010 Nov;56(5):481-9
6. Lyon RM. Significant modification of traditional rapid sequence induction improves safety and effectiveness of pre-hospital trauma anaesthesia. Crit Care. 2015 Apr 1;19:134.

7. Bruder EA. Single induction dose of etomidate versus other induction agents for endotracheal intubation in critically ill patients. Cochrane Database Syst Rev. 2015 Jan 8;1:CD010225.
8. Mace SE. Challenges and advances in intubation: rapid sequence intubation. Emerg Med Clin North Am. 2008 Nov;26(4):1043-68